



Grade 11/12 Math Circles Cardinality II - Problem Set

March 30 - April 6, 2022

Zack Cramer

1. Answer the following questions involving the cardinality of power sets.

(a) Let $A = \{a, b, c, d\}$. List the elements of $\mathcal{P}(A)$.

(b) If $|\mathcal{P}(A)| = 8192$, what is $|A|$?

(c) Does there exist a set A such that $|\mathcal{P}(A)| = 100$? Explain.

(d) If $|\mathcal{P}(\mathcal{P}(A))| = 2$, what can be said about A ?

(e) If $|\mathcal{P}(\mathcal{P}(A))|$ is less than 4 billion, what is the largest possible value of $|A|$?

2. Categorize the following sets based on their cardinality:

$$\mathbb{Z}, \mathcal{P}(\mathbb{Z}_{11}), \mathbb{R} \times \mathbb{R} \times \mathbb{R}, \mathbb{Z}_2 \times \mathbb{Z}_8, \mathbb{N} \times \mathbb{Q}, \mathcal{P}(\mathbb{N}), \mathcal{P}(\mathbb{R}), [0, 1], \mathcal{P}(\{0, 1\} \times \{0, 1\})$$

3. Let A, B, C, D be sets. Show that if $|A| = |C|$ and $|B| = |D|$, then $|A \times B| = |C \times D|$.

4. We have seen that a Cartesian product of finitely many countable sets is countable. That is, if A_1, A_2, \dots, A_n are countable, then so is $A_1 \times A_2 \times \dots \times A_n$.

Is the same true for a *countably infinite* collection of countable sets? That is, if A_1, A_2, A_3, \dots are countable sets, must $A_1 \times A_2 \times A_3 \times \dots$ be countable as well?

Hint: Let $A = \{0, 1, 2, \dots, 9\}$. Is the Cartesian product $A \times A \times A \times \dots$ countable? Think about decimal expansions.

5. Prove that at any point (a, b) in the xy -plane, there is a circle centred at (a, b) that does not pass through any points of the form (p, q) where p and q are rational.

Hint: Compare the number of circles one can draw at an arbitrary point (a, b) with the number of points (p, q) where p and q are rational.



6. Recall from one of your earlier Math Circles lessons that a real number α is said to be **algebraic** if there is a polynomial

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0,$$

where a_0, a_1, \dots, a_n are rational numbers, such that $p(\alpha) = 0$. For instance, $\sqrt{5}$ is algebraic, since $p(x) = x^2 - 5$ is a polynomial with rational coefficients and $p(\sqrt{5}) = 0$. If no such polynomial exists, α is said to be **transcendental**.

In this exercise, you will determine the cardinality of the set of algebraic numbers and the set of transcendental numbers.

- (a) Let \mathbb{P}_n be the set of all polynomials of degree n with rational coefficients. For instance, \mathbb{P}_2 is the set of all polynomials of the form

$$p(x) = a_2 x^2 + a_1 x + a_0, \text{ where } a_0, a_1, a_2 \in \mathbb{Q}.$$

Show that \mathbb{P}_n is countable by exhibiting a bijection

$$f : \mathbb{Q}^{n+1} \longrightarrow \mathbb{P}_n.$$

- (b) Let \mathbb{P} be the set of all polynomials with rational coefficients. Show that \mathbb{P} is countable.

Hint: Proposition 1 from the notes.

- (c) Let \mathbb{A} denote the set of algebraic real numbers. Using part (b), as well as the fact that a polynomial of degree n has at most n real roots, show that \mathbb{A} is countable.
- (d) Let \mathbb{T} denote the set of all transcendental real numbers. Show that \mathbb{T} is uncountable.¹

¹This shows that although it's tough to write down specific examples of transcendental numbers, most real numbers are, in fact, transcendental!